Rabbit anti-Human Cytokeratin 8 Polyclonal Antibody | anti-KRT8 antibody

Anti-Cytokeratin 8 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259783	**Rabbit**	**Human**	**Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot**

PRODUCT SYNONYMS

Cytokeratin 8; Polyclonal Antibody; Anti-Cytokeratin 8 Antibody; CYK8; Keratin; type II cytoskeletal 8; Cytokeratin-8; CK-8; Keratin-8; K8; Type-II keratin Kb8; anti-KRT8 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Cytokeratin 8 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	483
Applicable Applications for anti-KRT8 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Cytokeratin 8. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES

IF (Immunofluorescence)
Immunofluorescent analysis of Cytokeratin 8 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

IHC (Immunohiostchemistry)
Immunohistochemical analysis of Cytokeratin 8 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

WB (Western Blot)
Western blot analysis of Cytokeratin 8 expression in HeLa (A), A431 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-KRT8 antibody	Rabbit polyclonal antibody to Cytokeratin 8

NCBI AND UNIPROT INFORMATION

NCBI GI #	372466577
NCBI GeneID	3856
NCBI Accession #	NP_001243222.1
NCBI GenBank Nucleotide #	NM_001256293.1
UniProt Accession #	P05787; Q14099; Q14716; Q14717; Q53GJ0; Q6DHW5; Q6GMY0; Q6P4C7; Q96J60; A8K4H3; B0AZN5; F8VXB4
Molecular Weight	56,608 Da
NCBI Official Full Name	keratin, type II cytoskeletal 8 isoform 2
NCBI Official Synonym Full Names	keratin 8, type II
NCBI Official Symbol	KRT8
NCBI Official Synonym Symbols	K8; KO; CK8; CK-8; CYK8; K2C8; CARD2
NCBI Protein Information	keratin, type II cytoskeletal 8
UniProt Protein Name	Keratin, type II cytoskeletal 8
UniProt Gene Name	**KRT8**
UniProt Synonym Gene Names	CYK8; CK-8; K8
UniProt Entry Name	K2C8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/polyclonal/cytokeratin-8-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/259783