

Rabbit anti-Human Cytokeratin 18 Polyclonal Antibody | anti-KRT18 antibody

Anti-Cytokeratin 18 (pS33) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259788	Rabbit	Human	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Cytokeratin 18; Polyclonal Antibody; Anti-Cytokeratin 18 (pS33) Phospho Antibody; CYK18; Keratin; type I cytoskeletal 18; Cell proliferation-inducing gene 46 protein; Cytokeratin-18; CK-18; Keratin-18; K18; anti-KRT18 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Cytokeratin 18 (pS33) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	430
Applicable Applications for anti-KRT18 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Cytokeratin 18. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Cytokeratin 18 (pS33) staining in A431 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Cytokeratin 18 (pS33) staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
 Western blot analysis of Cytokeratin 18 (pS33) expression in A431 (A), HeLa (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-KRT18 antibody	Rabbit polyclonal antibody to Cytokeratin 18 (pS33)
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557888
NCBI GeneID	3875
NCBI Accession #	NP_000215.1
NCBI GenBank Nucleotide #	NM_000224.2
UniProt Accession #	P05783; Q53G38; Q5U0N8; Q9BW26
Molecular Weight	48,058 Da
NCBI Official Full Name	keratin, type I cytoskeletal 18
NCBI Official Synonym Full Names	keratin 18, type I
NCBI Official Symbol	KRT18
NCBI Official Synonym Symbols	K18; CK-18; CYK18
NCBI Protein Information	keratin, type I cytoskeletal 18
UniProt Protein Name	Keratin, type I cytoskeletal 18
UniProt Gene Name	KRT18
UniProt Synonym Gene Names	CYK18; CK-18; K18
UniProt Entry Name	K1C18_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.