

Rabbit MEF2C Polyclonal Antibody | anti-MEF2C antibody

Anti-MEF2C Antibody

CATALOG # AAA259823	HOST Rabbit	REACTIVITY Human, Mouse, Porcine, Zebrafish	APPLICATIONS Chromatin Immunoprecipitation, Immunoprecipitation, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	---	--


PRODUCT SYNONYMS

MEF2C; Polyclonal Antibody; Anti-MEF2C Antibody; Myocyte-specific enhancer factor 2C; anti-MEF2C antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Porcine, Zebrafish
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of MEF2C protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	463
Applicable Applications for anti-MEF2C antibody	ChIP (Chromatin immunoprecipitation), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human MEF2C. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of MEF2C staining in PC12 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of MEF2C staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MEF2C expression in HeLa (A), NIH3T3 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-MEF2C antibody	Rabbit polyclonal antibody to MEF2C
---	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	196114945
NCBI GeneID	4208
NCBI Accession #	NP_001124477.1
NCBI GenBank Nucleotide #	NM_001131005.2
UniProt Accession #	Q06413; C9JMZ0; D7F7N5; F8W7V7
Molecular Weight	50,242 Da
NCBI Official Full Name	myocyte-specific enhancer factor 2C isoform 2
NCBI Official Synonym Full Names	myocyte enhancer factor 2C
NCBI Official Symbol	MEF2C
NCBI Official Synonym Symbols	DEL5q14.3; C5DELq14.3
NCBI Protein Information	myocyte-specific enhancer factor 2C
UniProt Protein Name	Myocyte-specific enhancer factor 2C
UniProt Gene Name	MEF2C
UniProt Entry Name	MEF2C_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mef2c-antibody-chromatin-immunoprecipitation-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/259823>