

Rabbit Ataxin 1 Polyclonal Antibody | anti-ATXN1 antibody

Anti-Ataxin 1 (pS775) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259839	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Ataxin 1; Polyclonal Antibody; Anti-Ataxin 1 (pS775) Phospho Antibody; ATX1; SCA1; Ataxin-1; Spinocerebellar ataxia type 1 protein; anti-ATXN1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Ataxin 1 (pS775) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	815
Applicable Applications for anti-ATXN1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Ataxin 1 (pS775). The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Ataxin 1 (pS775) staining in HEK293 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Ataxin 1 (pS775) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
 Western blot analysis of Ataxin 1 (pS775) expression in HEK293T (A), mouse brain (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-ATXN1 antibody	Rabbit polyclonal antibody to Ataxin 1 (pS775)
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	51479158
NCBI GeneID	6310
NCBI Accession #	NP_000323.2
NCBI GenBank Nucleotide #	NM_000332.3
UniProt Accession #	P54253; Q17S02; Q9UJG2; Q9Y4J1
Molecular Weight	86,923 Da
NCBI Official Full Name	ataxin-1
NCBI Official Synonym Full Names	ataxin 1
NCBI Official Symbol	ATXN1
NCBI Official Synonym Symbols	ATX1; SCA1; D6S504E
NCBI Protein Information	ataxin-1
UniProt Protein Name	Ataxin-1
UniProt Gene Name	ATXN1
UniProt Synonym Gene Names	ATX1; SCA1
UniProt Entry Name	ATX1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/ataxin-1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/259839>