

anti-Human, Rat MUC16 Polyclonal Antibody | anti-MUC16 antibody

Anti-MUC16 Antibody

CATALOG # AAA259918	REACTIVITY Human, Rat	APPLICATIONS Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot
-------------------------------	---------------------------------	--

PRODUCT SYNONYMS

MUC16; Polyclonal Antibody; Anti-MUC16 Antibody; CA125; Mucin-16; MUC-16; Ovarian cancer-related tumor marker CA125; CA-125; Ovarian carcinoma antigen CA125; anti-MUC16 antibody

PRODUCT SPECIFICATIONS

Reactivity	Human, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of MUC16 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	14507
Applicable Applications for anti-MUC16 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Source	Rabbit
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human MUC16. The exact sequence is proprietary.
Preparation and Storage	Upon receipt aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of MUC16 staining in HuvEc cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of MUC16 staining in human bladder cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MUC16 expression in MCF7 (A), H9C2 (B), PC12 (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-MUC16 antibody	Rabbit polyclonal antibody to MUC16
---	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	83367077
NCBI GeneID	94025
NCBI Accession #	NP_078966.2
NCBI GenBank Nucleotide #	NM_024690.2
UniProt Accession #	Q8WXI7; Q6ZQW5; Q96RK2
NCBI Official Full Name	mucin-16
NCBI Official Synonym Full Names	mucin 16, cell surface associated
NCBI Official Symbol	MUC16
NCBI Official Synonym Symbols	CA125
NCBI Protein Information	mucin-16
UniProt Protein Name	Mucin-16
UniProt Gene Name	MUC16
UniProt Synonym Gene Names	CA125; MUC-16; CA-125
UniProt Entry Name	MUC16_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/muc16-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/259918>