

Rabbit Tyrosine Hydroxylase Polyclonal Antibody | anti-TH antibody

Anti-Tyrosine Hydroxylase Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260015	Rabbit	Human, Mouse, Rat, Chicken, Dog	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Tyrosine Hydroxylase; Polyclonal Antibody; Anti-Tyrosine Hydroxylase Antibody; TYH; Tyrosine 3-monooxygenase; Tyrosine 3-hydroxylase; TH; anti-TH antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Chicken, Dog
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Tyrosine Hydroxylase protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	497
Applicable Applications for anti-TH antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Tyrosine Hydroxylase. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Tyrosine Hydroxylase staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Tyrosine Hydroxylase staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Tyrosine Hydroxylase expression in A549 (A), NIH3T3 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-TH antibody	Rabbit polyclonal antibody to Tyrosine Hydroxylase
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	88900503
NCBI GeneID	7054
NCBI Accession #	NP_000351.2
NCBI GenBank Nucleotide #	NM_000360.3
UniProt Accession #	P07101; Q0PWM2; Q0PWM3; Q15585; Q15588; Q15589; Q2M3B4; B7ZL70; B7ZL73
Molecular Weight	44,898 Da
NCBI Official Full Name	tyrosine 3-monooxygenase isoform b
NCBI Official Synonym Full Names	tyrosine hydroxylase
NCBI Official Symbol	TH
NCBI Official Synonym Symbols	TYH; DYT14; DYT5b
NCBI Protein Information	tyrosine 3-monooxygenase
UniProt Protein Name	Tyrosine 3-monooxygenase
UniProt Gene Name	TH
UniProt Synonym Gene Names	TYH; TH
UniProt Entry Name	TY3H_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tyrosine-hydroxylase-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/260015>