

Rabbit PAK1 Polyclonal Antibody | anti-PAK1 antibody

Anti-PAK1 (pT212) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260057	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PAK1; Polyclonal Antibody; Anti-PAK1 (pT212) Phospho Antibody; Serine/threonine-protein kinase PAK 1; Alpha-PAK; p21-activated kinase 1; PAK-1; p65-PAK; anti-PAK1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of PAK1 (pT212) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	553
Applicable Applications for anti-PAK1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human PAK1 (pT212). The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of PAK1 (pT212) staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of PAK1 (pT212) staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of PAK1 (pT212) expression in HCT116 (A) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-PAK1 antibody

Rabbit polyclonal antibody to PAK1 (pT212)

NCBI AND UNIPROT INFORMATION

NCBI GI #	190886457
NCBI GeneID	5058
NCBI Accession #	NP_001122092.1
NCBI GenBank Nucleotide #	NM_001128620.1
UniProt Accession #	Q13153; O75561; Q13567; Q32M53; Q32M54; Q86W79
Molecular Weight	61,632 Da
NCBI Official Full Name	serine/threonine-protein kinase PAK 1 isoform 1
NCBI Official Synonym Full Names	p21 protein (Cdc42/Rac)-activated kinase 1
NCBI Official Symbol	PAK1
NCBI Official Synonym Symbols	PAKalpha
NCBI Protein Information	serine/threonine-protein kinase PAK 1
UniProt Protein Name	Serine/threonine-protein kinase PAK 1
UniProt Gene Name	PAK1
UniProt Synonym Gene Names	PAK-1
UniProt Entry Name	PAK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/pak1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/260057>