

Rabbit CaMK1 alpha Polyclonal Antibody | anti-CAMK1 antibody

Anti-CaMK1 alpha (pT177) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260161	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CaMK1 alpha; Polyclonal Antibody; Anti-CaMK1 alpha (pT177) Phospho Antibody; Calcium/calmodulin-dependent protein kinase type 1; CaM kinase I; CaM-KI; CaM kinase I alpha; CaMKI-alpha; anti-CAMK1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of CaMK1 alpha (pT177) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	370
Applicable Applications for anti-CAMK1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CaMK1 alpha. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of CaMK1 alpha (pT177) staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of CaMK1 alpha (pT177) staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
Western blot analysis of CaMK1 alpha (pT177) expression in Jurkat Insulin-treated (A) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-CAMK1 antibody	Rabbit polyclonal antibody to CaMK1 alpha (pT177)
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502553
NCBI GeneID	8536
NCBI Accession #	NP_003647.1
NCBI GenBank Nucleotide #	NM_003656.4
UniProt Accession #	Q14012; Q3KPF6
Molecular Weight	41,337 Da
NCBI Official Full Name	calcium/calmodulin-dependent protein kinase type 1
NCBI Official Synonym Full Names	calcium/calmodulin-dependent protein kinase I
NCBI Official Symbol	CAMK1
NCBI Official Synonym Symbols	CAMKI
NCBI Protein Information	calcium/calmodulin-dependent protein kinase type 1
UniProt Protein Name	Calcium/calmodulin-dependent protein kinase type 1
UniProt Gene Name	CAMK1
UniProt Synonym Gene Names	CaM-KI; CaMKI-alpha
UniProt Entry Name	KCC1A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/camk1-alpha-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/260161>