

Rabbit RET Polyclonal Antibody | anti-RET antibody

Anti-RET Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260236	Rabbit	Human, Mouse, Rat, Monkey	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

RET; Polyclonal Antibody; Anti-RET Antibody; CDHF12; CDHR16; PTC; RET51; Proto-oncogene tyrosine-protein kinase receptor Ret; Cadherin family member 12; Proto-oncogene c-Ret; anti-RET antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of RET protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	1072
Applicable Applications for anti-RET antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human RET. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of RET staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of RET staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of RET expression in K562 (A) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-RET antibody

Rabbit polyclonal antibody to RET

NCBI AND UNIPROT INFORMATION

NCBI GI #	10862701
NCBI GeneID	5979
NCBI Accession #	NP_065681.1
NCBI GenBank Nucleotide #	NM_020630.4
UniProt Accession #	P07949; Q15250; Q9BTB0; Q9H4A2; A8K6Z2
Molecular Weight	119,847 Da
NCBI Official Full Name	proto-oncogene tyrosine-protein kinase receptor Ret isoform c
NCBI Official Synonym Full Names	ret proto-oncogene
NCBI Official Symbol	RET
NCBI Official Synonym Symbols	PTC; MTC1; HSCR1; MEN2A; MEN2B; RET51; CDHF12; CDHR16; RET-ELE1
NCBI Protein Information	proto-oncogene tyrosine-protein kinase receptor Ret
UniProt Protein Name	Proto-oncogene tyrosine-protein kinase receptor Ret
UniProt Gene Name	RET
UniProt Synonym Gene Names	CDHF12; CDHR16; PTC; RET51
UniProt Entry Name	RET_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/ret-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/260236>