

Rabbit Claudin 3 Polyclonal Antibody | anti-CLDN3 antibody

Anti-Claudin 3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260277	Rabbit	Human, Bovine	Immunoprecipitation, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Claudin 3; Polyclonal Antibody; Anti-Claudin 3 Antibody; C7orf1; CPETR2; Claudin-3; Clostridium perfringens enterotoxin receptor 2; CPE-R 2; CPE-receptor 2; Rat ventral prostate.1 protein homolog; hRVP1; anti-CLDN3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Bovine
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Claudin 3 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	220
Applicable Applications for anti-CLDN3 antibody	IP (Immunoprecipitation), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Claudin 3. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Claudin 3 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Claudin 3 staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Claudin 3 expression in SW480 (A), LOVO (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-CLDN3 antibody

Rabbit polyclonal antibody to Claudin 3

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502875
NCBI GeneID	1365
NCBI Accession #	NP_001297.1
NCBI GenBank Nucleotide #	NM_001306.3
UniProt Accession #	O15551
Molecular Weight	23,319 Da
NCBI Official Full Name	claudin-3
NCBI Official Synonym Full Names	claudin 3
NCBI Official Symbol	CLDN3
NCBI Official Synonym Symbols	RVP1; HRVP1; C7orf1; CPE-R2; CPETR2
NCBI Protein Information	claudin-3
UniProt Protein Name	Claudin-3
UniProt Gene Name	CLDN3
UniProt Synonym Gene Names	C7orf1; CPETR2; CPE-R 2; CPE-receptor 2; hRVP1
UniProt Entry Name	CLD3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/claudin-3-antibody-immunoprecipitation-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/260277>