

Histone Deacetylase 6 Polyclonal Antibody | anti-HDAC6 antibody

Anti-Histone Deacetylase 6 (pS22) Phospho Antibody

CATALOG #	REACTIVITY	APPLICATIONS
AAA260329	Human, Monkey	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Histone Deacetylase 6; Polyclonal Antibody; Anti-Histone Deacetylase 6 (pS22) Phospho Antibody; KIAA0901; Histone deacetylase 6; HD6; anti-HDAC6 antibody

PRODUCT SPECIFICATIONS


Reactivity	Human, Monkey
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Histone Deacetylase 6 (pS22) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	1215
Applicable Applications for anti-HDAC6 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Source	Rabbit
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Histone Deacetylase 6 (pS22). The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Histone Deacetylase 6 (pS22) staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Histone Deacetylase 6 (pS22) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Histone Deacetylase 6 (pS22) expression in HepG2 (A), HeLa (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-HDAC6 antibody	Rabbit polyclonal antibody to Histone Deacetylase 6 (pS22)
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	13128864
NCBI GeneID	10013
NCBI Accession #	NP_006035.2
NCBI GenBank Nucleotide #	NM_006044.2
UniProt Accession #	Q9UBN7; O94975; Q6NT75; Q7L3E5; Q96CY0
Molecular Weight	114,361 Da
NCBI Official Full Name	histone deacetylase 6
NCBI Official Synonym Full Names	histone deacetylase 6
NCBI Official Symbol	HDAC6
NCBI Official Synonym Symbols	HD6; JM21; CPBHM; PPP1R90
NCBI Protein Information	histone deacetylase 6
UniProt Protein Name	Histone deacetylase 6
UniProt Gene Name	HDAC6
UniProt Synonym Gene Names	KIAA0901; HD6
UniProt Entry Name	HDAC6_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/histone-deacetylase-6-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/260329>