

Mouse anti-Human p53 Monoclonal Antibody | anti-TP53 antibody

Anti-p53 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260416	Mouse	Human	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

p53; Monoclonal Antibody; Anti-p53 Antibody; P53; Cellular tumor antigen p53; Antigen NY-CO-13; Phosphoprotein p53; Tumor suppressor p53; anti-TP53 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Specificity	Recognizes endogenous levels of p53 protein.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	393
Applicable Applications for anti-TP53 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence of human p53. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of p53 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a FITC-conjugated secondary antibody (green) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of p53 staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of p53 expression in 293T (A) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-TP53 antibody	Mouse monoclonal to p53
--	-------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	269849759
NCBI GeneID	7157
UniProt Accession #	P04637; Q15086; Q15087; Q15088; Q16535; Q16807; Q16808; Q16809; Q16810; Q16811; Q16848; Q2XN98
Molecular Weight	24,401 Da
NCBI Official Full Name	Cellular tumor antigen p53
NCBI Official Synonym Full Names	tumor protein p53
NCBI Official Symbol	TP53
NCBI Official Synonym Symbols	P53; BCC7; LFS1; TRP53
NCBI Protein Information	cellular tumor antigen p53
UniProt Protein Name	Cellular tumor antigen p53
UniProt Gene Name	TP53
UniProt Synonym Gene Names	P53
UniProt Entry Name	P53_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/p53-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/260416>