

Mouse HSP70 Monoclonal Antibody | anti-HSPA1L antibody

Anti-HSP70 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260422	Mouse	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

HSP70; Monoclonal Antibody; Anti-HSP70 Antibody; HSPA1L; Heat shock 70 kDa protein 1-like; Heat shock 70 kDa protein 1L; Heat shock 70 kDa protein 1-Hom; HSP70-Hom; HSPA1A; HSPA1; HSPA1B; Heat shock 70 kDa protein 1A/1B; Heat shock 70 kDa protein 1/2; HSP70-1/HSP70-2; HSP70.1/HSP70.2; anti-HSPA1L antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Specificity	Recognizes endogenous levels of HSP70 protein.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	641
Applicable Applications for anti-HSPA1L antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence of human HSP70. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of HSP70 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a FITC-conjugated secondary antibody (green) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of HSP70 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of HSP70 expression in Hela (A), mouse brain (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-HSPA1L antibody	Mouse monoclonal to HSP70
--	---------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	23831140
NCBI GeneID	3305
UniProt Accession #	P34931; O75634; Q2HXR3; Q8NE72; Q96QC9; Q9UQM1; A6NNB0; B0UXW8
Molecular Weight	70,375 Da
NCBI Official Full Name	Heat shock 70 kDa protein 1-like
NCBI Official Synonym Full Names	heat shock protein family A (Hsp70) member 1 like
NCBI Official Symbol	HSPA1L
NCBI Official Synonym Symbols	HSP70T; hum70t; HSP70-1L; HSP70-HOM
NCBI Protein Information	heat shock 70 kDa protein 1-like
UniProt Protein Name	Heat shock 70 kDa protein 1-like
UniProt Gene Name	HSPA1L
UniProt Synonym Gene Names	Heat shock 70 kDa protein 1L; HSP70-Hom
UniProt Entry Name	HS71L_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/hsp70-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/260422>