

Mouse Myc-tag Monoclonal Antibody | anti-Myc antibody

Anti-Myc-tag Antibody

CATALOG #	HOST	APPLICATIONS
AAA260424	Mouse	Immunoprecipitation, Immunocytochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

Myc-tag; Monoclonal Antibody; Anti-Myc-tag Antibody; anti-Myc antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Clonality	Monoclonal
Specificity	Recognizes Myc-tag fusion proteins.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	11
Applicable Applications for anti-Myc antibody	IP (Immunoprecipitation), ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence of Myc-tag. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IP (Immunoprecipitation)

Immunoprecipitation of V5-tagged protein from HEK293T cells transfected with vector overexpressing V5 tag, using anti-V5-tag Antibody.



IF (Immunofluorescence)

Immunofluorescent analysis of V5-tag staining in 293T cells transfected with a V5-tag protein. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



WB (Western Blot)

Western blot analysis of over-expressed V5-tagged protein in 293T cell lysate.

ADDITIONAL INFORMATION

Related Product Information for anti-Myc antibody

Mouse monoclonal to Myc-tag

NCBI AND UNIPROT INFORMATION

NCBI GI #

985015742

NCBI Official Full Name

MYC-tag, partial

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/myc-tag-antibody-immunoprecipitation-immunocytochemistry-immunofluorescence-western-blot-elisa/260424>