

Rabbit anti-Human MCL1 Polyclonal Antibody | anti-MCL1 antibody

Anti-MCL1 (pS159) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260561	Rabbit	Human	Immunocytochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

MCL1; Polyclonal Antibody; Anti-MCL1 (pS159) Phospho Antibody; BCL2L3; Induced myeloid leukemia cell differentiation protein Mcl-1; Bcl-2-like protein 3; Bcl2-L-3; Bcl-2-related protein EAT/mcl1; mcl1/EAT; anti-MCL1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of MCL1 (pS159) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	350
Applicable Applications for anti-MCL1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human MCL1. The exact sequence is proprietary.
Preparation and Storage	Shipped and store at 4 degree C for one year. Do not freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of MCL1 (pS159) staining in K562 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



WB (Western Blot)

Western blot analysis of MCL1 (pS159) expression in K562 (A), human liver (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-MCL1 antibody	Rabbit polyclonal antibody to MCL1 (pS159)
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	83304396
NCBI GeneID	4170
UniProt Accession #	Q07820; Q9HD91; Q9NRQ3; Q9NRQ4; Q9UHR7; Q9UHR8; Q9UHR9; Q9UNJ1; B2R6B2; D3DV03; D3DV04
Molecular Weight	28,662 Da
NCBI Official Full Name	Induced myeloid leukemia cell differentiation protein Mcl-1
NCBI Official Synonym Full Names	myeloid cell leukemia 1
NCBI Official Symbol	MCL1
NCBI Official Synonym Symbols	TM; EAT; MCL1L; MCL1S; Mcl-1; BCL2L3; MCL1-ES; bcl2-L-3; mcl1/EAT
NCBI Protein Information	induced myeloid leukemia cell differentiation protein Mcl-1
UniProt Protein Name	Induced myeloid leukemia cell differentiation protein Mcl-1
UniProt Gene Name	MCL1
UniProt Synonym Gene Names	BCL2L3; Bcl2-L-3
UniProt Entry Name	MCL1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.