

Rabbit Ezrin/Radixin/Moesin Polyclonal Antibody | anti-MSN antibody

Anti-Ezrin/Radixin/Moesin (pT567/564/558) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260603	Rabbit	Human, Mouse, Rat, Bovine, Monkey, Pig, Zebrafish	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Ezrin/Radixin/Moesin; Polyclonal Antibody; Anti-Ezrin/Radixin/Moesin (pT567/564/558) Phospho Antibody; MSN; Moesin; Membrane-organizing extension spike protein; RDX; Radixin; EZR; VIL2; Ezrin; Cytovillin; Villin-2; p81; anti-MSN antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Monkey, Pig, Zebrafish
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Ezrin/Radixin/Moesin (pT567/564/558) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	577
Applicable Applications for anti-MSN antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Ezrin/Radixin/Moesin. The exact sequence is proprietary.
Preparation and Storage	Shipped and store at 4 degree C for one year. Do not freeze.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Ezrin/Radixin/Moesin (pT567/564/558) staining in SKOV3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Ezrin/Radixin/Moesin (pT567/564/558) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
 Western blot analysis of Ezrin/Radixin/Moesin (pT567/564/558) expression in SKOV3 (A), NIH3T3 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-MSN antibody	Rabbit polyclonal antibody to Ezrin/Radixin/Moesin (pT567/564/558)
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	127234
NCBI GeneID	4478
UniProt Accession #	P26038; P35241; P15311; P26041; P26043; P26040; O35763; P31977
Molecular Weight	67,820 Da
NCBI Official Full Name	Moesin
NCBI Official Synonym Full Names	moesin
NCBI Official Symbol	MSN
NCBI Official Synonym Symbols	HEL70
NCBI Protein Information	moesin
UniProt Protein Name	Moesin
UniProt Gene Name	MSN
UniProt Entry Name	MOES_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/ezrin-radixin-moesin-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/260603>