

Rabbit TRK A Polyclonal Antibody | anti-NTRK1 antibody

Anti-TRK A (pY496) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260620	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

TRK A; Polyclonal Antibody; Anti-TRK A (pY496) Phospho Antibody; MTC; TRK; TRKA; High affinity nerve growth factor receptor; Neurotrophic tyrosine kinase receptor type 1; TRK1-transforming tyrosine kinase protein; Tropomyosin-related kinase A; Tyrosine kinase receptor; Tyrosine kinase receptor A; Trk-A; gp140trk; p140-TrkA; anti-NTRK1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of TRK A (pY496) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	796
Applicable Applications for anti-NTRK1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TRK A. The exact sequence is proprietary.
Preparation and Storage	Shipped and store at 4 degree C for one year. Do not freeze.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of TRK A (pY496) staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of TRK A (pY496) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of TRK A (pY496) expression in mouse brain (A), rat brain (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-NTRK1 antibody	Rabbit polyclonal antibody to TRK A (pY496)
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	94730402
NCBI GeneID	4914
UniProt Accession #	P04629; P08119; Q15655; Q15656; Q5D056; Q5VZS2; Q7Z5C3; Q9UIU7; B2R6T5; B7ZM34
Molecular Weight	77,145 Da
NCBI Official Full Name	High affinity nerve growth factor receptor
NCBI Official Synonym Full Names	neurotrophic receptor tyrosine kinase 1
NCBI Official Symbol	NTRK1
NCBI Official Synonym Symbols	MTC; TRK; TRK1; TRKA; Trk-A; p140-TrkA
NCBI Protein Information	high affinity nerve growth factor receptor
UniProt Protein Name	High affinity nerve growth factor receptor
UniProt Gene Name	NTRK1
UniProt Synonym Gene Names	MTC; TRK; TRKA; Trk-A
UniProt Entry Name	NTRK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/trk-a-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/260620>