

Rabbit AMPK alpha 1 Polyclonal Antibody | anti-PRKAA1 antibody

Anti-AMPK alpha 1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260631	Rabbit	Human, Mouse, Rat, Chicken	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

AMPK alpha 1; Polyclonal Antibody; Anti-AMPK alpha 1 Antibody; AMPK1; 5'-AMP-activated protein kinase catalytic subunit alpha-1; AMPK subunit alpha-1; Acetyl-CoA carboxylase kinase; ACACA kinase; Hydroxymethylglutaryl-CoA reductase kinase; HMGCR kinase; Tau-protein kinase PRKAA1; anti-PRKAA1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Chicken
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of AMPK alpha 1 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	559
Applicable Applications for anti-PRKAA1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human AMPK alpha 1. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of AMPK alpha 1 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of AMPK alpha 1 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of AMPK alpha 1 expression in HeLa (A), K562 (B), NIH3T3 (C), H9C2 (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-PRKAA1 antibody	Rabbit polyclonal antibody to AMPK alpha 1
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	254763436
NCBI GeneID	5562
UniProt Accession #	Q13131; O00286; Q5D0E1; Q86VS1; Q9UNQ4; A8MTQ6; B2R7E1
Molecular Weight	65,523 Da
NCBI Official Full Name	5'-AMP-activated protein kinase catalytic subunit alpha-1
NCBI Official Synonym Full Names	protein kinase AMP-activated catalytic subunit alpha 1
NCBI Official Symbol	PRKAA1
NCBI Official Synonym Symbols	AMPK; AMPKa1
NCBI Protein Information	5'-AMP-activated protein kinase catalytic subunit alpha-1
UniProt Protein Name	5'-AMP-activated protein kinase catalytic subunit alpha-1
UniProt Gene Name	PRKAA1
UniProt Synonym Gene Names	AMPK1; HMGCR kinase
UniProt Entry Name	AAPK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.