

Rabbit c-Met Polyclonal Antibody | anti-MET antibody

Anti-c-Met Antibody

CATALOG # AAA260676	HOST Rabbit	REACTIVITY Human, Mouse, Rat, Rabbit, Sheep	APPLICATIONS Immunocytochemistry, Immunofluorescence, Western Blot
-------------------------------	-----------------------	---	--

PRODUCT SYNONYMS

c-Met; Polyclonal Antibody; Anti-c-Met Antibody; Hepatocyte growth factor receptor; HGF receptor; HGF/SF receptor; Proto-oncogene c-Met; Scatter factor receptor; SF receptor; Tyrosine-protein kinase Met; anti-MET antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Rabbit, Sheep
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of c-Met protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	1390
Applicable Applications for anti-MET antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human c-Met. The exact sequence is proprietary.
Preparation and Storage	Shipped and store at 4 degree C for one year. Do not freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of c-Met staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



WB (Western Blot)

Western blot analysis of c-Met expression in HeLa (A), A431 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-MET antibody	Rabbit polyclonal antibody to c-Met
---	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	251757497
-----------	-----------

NCBI GeneID	4233
UniProt Accession #	P08581; O60366; Q12875; Q9UDX7; Q9UPL8; A1L467; B5A932; E7EQ94
Molecular Weight	85,745 Da
NCBI Official Full Name	Hepatocyte growth factor receptor
NCBI Official Synonym Full Names	MET proto-oncogene, receptor tyrosine kinase
NCBI Official Symbol	MET
NCBI Official Synonym Symbols	HGFR; AUTS9; RCCP2; c-Met; DFNB97
NCBI Protein Information	hepatocyte growth factor receptor
UniProt Protein Name	Hepatocyte growth factor receptor
UniProt Gene Name	MET
UniProt Synonym Gene Names	HGF receptor; SF receptor
UniProt Entry Name	MET_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/c-met-antibody-immunocytochemistry-immunofluorescence-western-blot-elisa/260676>