

Rabbit CD226 Polyclonal Antibody | anti-CD226 antibody

Anti-CD226 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260747	Rabbit	Human, Mouse, Monkey	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

CD226; Polyclonal Antibody; Anti-CD226 Antibody; DNAM1; CD226 antigen; DNAX accessory molecule 1; DNAM-1; anti-CD226 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Monkey
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of CD226 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	336
Applicable Applications for anti-CD226 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CD226. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of CD226 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of CD226 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of CD226 expression in HEK293T (A), mouse brain (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-CD226 antibody	Rabbit polyclonal antibody to CD226
---	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	317373338
NCBI GeneID	10666
UniProt Accession #	Q15762; B2R818
Molecular Weight	38,614 Da
NCBI Official Full Name	CD226 antigen
NCBI Official Synonym Full Names	CD226 molecule
NCBI Official Symbol	CD226
NCBI Official Synonym Symbols	PTA1; DNAM1; DNAM-1; TLISA1
NCBI Protein Information	CD226 antigen
UniProt Protein Name	CD226 antigen
UniProt Gene Name	CD226
UniProt Synonym Gene Names	DNAM1; DNAM-1
UniProt Entry Name	CD226_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd226-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/260747>