

Rabbit MGCRCAGAP Polyclonal Antibody | anti-RACGAP1 antibody

Anti-MGCRCAGAP Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260816	Rabbit	Human, Mouse, Monkey	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MGCRCAGAP; Polyclonal Antibody; Anti-MGCRCAGAP Antibody; KIAA1478; Rac GTPase-activating protein 1; Male germ cell RacGap; MgcRacGAP; Protein CYK4 homolog; CYK4; HsCYK-4; anti-RACGAP1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Monkey
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of MGCRCAGAP protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	632
Applicable Applications for anti-RACGAP1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human MGCRCAGAP. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of MGCRCAGAP staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of MGCRCAGAP staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MGCRCACGAP expression in A431 (A), Raji (B), COS7 (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-RACGAP1 antibody

Rabbit polyclonal antibody to MGCRCACGAP

NCBI AND UNIPROT INFORMATION

NCBI GI #	74762727
NCBI GeneID	29127
UniProt Accession #	Q9H0H5; Q6PJ26; Q9NWN2; Q9P250; Q9P2W2
Molecular Weight	71,027 Da
NCBI Official Full Name	Rac GTPase-activating protein 1
NCBI Official Synonym Full Names	Rac GTPase activating protein 1
NCBI Official Symbol	RACGAP1
NCBI Official Synonym Symbols	CYK4; ID-GAP; HsCYK-4; MgcRacGAP
NCBI Protein Information	rac GTPase-activating protein 1
UniProt Protein Name	Rac GTPase-activating protein 1
UniProt Gene Name	RACGAP1
UniProt Synonym Gene Names	MgcRacGAP; CYK4; HsCYK-4
UniProt Entry Name	RGAP1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mgcracgap-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/260816>