

Rabbit PPAR gamma Polyclonal Antibody | anti-PPARG antibody

Anti-PPAR gamma Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260931	Rabbit	Human, Mouse, Rat, Dog, Rabbit	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PPAR gamma; Polyclonal Antibody; Anti-PPAR gamma Antibody; NR1C3; Peroxisome proliferator-activated receptor gamma; PPAR-gamma; Nuclear receptor subfamily 1 group C member 3; anti-PPARG antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Dog, Rabbit
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of PPAR gamma protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	505
Applicable Applications for anti-PPARG antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human PPAR gamma. The exact sequence is proprietary.
Preparation and Storage	Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of PPAR gamma staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of PPAR gamma staining in human colon formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of PPAR gamma expression in Hela (A), rat ovary (B), mouse spleen (C), LO2 (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-PPARG antibody

Rabbit polyclonal antibody to PPAR gamma

NCBI AND UNIPROT INFORMATION

NCBI GI #	13432234
NCBI GeneID	5468
UniProt Accession #	P37231; O00684; O00710; O14515; Q0QJH8; Q15178; Q15179; Q15180; Q15832; Q86U60; A8K3G6; B5BUA1
NCBI Official Full Name	Peroxisome proliferator-activated receptor gamma
NCBI Official Synonym Full Names	peroxisome proliferator activated receptor gamma
NCBI Official Symbol	PPARG
NCBI Official Synonym Symbols	GLM1; CIMT1; NR1C3; PPARG1; PPARG2; PPARGgamma
NCBI Protein Information	peroxisome proliferator-activated receptor gamma
UniProt Protein Name	Peroxisome proliferator-activated receptor gamma
UniProt Gene Name	PPARG
UniProt Synonym Gene Names	NR1C3; PPAR-gamma
UniProt Entry Name	PPARG_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/ppar-gamma-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/260931>