

Rabbit Dyskerin Polyclonal Antibody | anti-DKC1 antibody

Anti-Dyskerin Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA260990	Rabbit	Human, Mouse, Monkey, Rabbit	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Dyskerin; Polyclonal Antibody; Anti-Dyskerin Antibody; NOLA4; H/ACA ribonucleoprotein complex subunit 4; CBF5 homolog; Nopp140-associated protein of 57 kDa; Nucleolar protein NAP57; Nucleolar protein family A member 4; snoRNP protein DKC1; anti-DKC1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Monkey, Rabbit
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Dyskerin protein.
Purity/Purification	The antibody was purified by affinity chromatography.
Form/Format	Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% Sodium Azide.
Sequence Length	514
Applicable Applications for anti-DKC1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Dyskerin. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Dyskerin staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Dyskerin staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Dyskerin expression in MCF7 (A), HepG2 (B), Jurkat (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-DKC1 antibody	Rabbit polyclonal antibody to Dyskerin
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	3913488
NCBI GeneID	1736
UniProt Accession #	O60832; O43845; Q96G67; Q9Y505; F5BSB3
Molecular Weight	47,603 Da
NCBI Official Full Name	H/ACA ribonucleoprotein complex subunit 4
NCBI Official Synonym Full Names	dyskerin pseudouridine synthase 1
NCBI Official Symbol	DKC1
NCBI Official Synonym Symbols	DKC; CBF5; DKCX; NAP57; NOLA4; XAP101
NCBI Protein Information	H/ACA ribonucleoprotein complex subunit 4
UniProt Protein Name	H/ACA ribonucleoprotein complex subunit 4
UniProt Gene Name	DKC1
UniProt Synonym Gene Names	NOLA4
UniProt Entry Name	DKC1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/dyskerin-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/260990>