

Rabbit Nucleophosmin Polyclonal Antibody | anti-NPM1 antibody

Anti-Nucleophosmin Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA261092	Rabbit	Human, Mouse, Rat, Bovine, Rabbit	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Nucleophosmin; Polyclonal Antibody; Anti-Nucleophosmin Antibody; NPM; Nucleolar phosphoprotein B23; Nucleolar protein NO38; Numatrin; anti-NPM1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Rabbit
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Nucleophosmin protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	294
Applicable Applications for anti-NPM1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Nucleophosmin. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Nucleophosmin staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Nucleophosmin staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Nucleophosmin expression in COLO205 (A), NIH3T3 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-NPM1 antibody

Rabbit polyclonal antibody to Nucleophosmin

NCBI AND UNIPROT INFORMATION

NCBI GI #	114762
NCBI GeneID	4869
UniProt Accession #	P06748; P08693; Q12826; Q13440; Q13441; Q14115; Q5EU94; Q5EU95; Q5EU96; A8K3N7; B5BU00; D3DQL6
Molecular Weight	28,400 Da
NCBI Official Full Name	Nucleophosmin
NCBI Official Synonym Full Names	nucleophosmin (nucleolar phosphoprotein B23, numatrin)
NCBI Official Symbol	NPM1
NCBI Official Synonym Symbols	B23; NPM
NCBI Protein Information	nucleophosmin
UniProt Protein Name	Nucleophosmin
UniProt Gene Name	NPM1
UniProt Synonym Gene Names	NPM; NPM
UniProt Entry Name	NPM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/nucleophosmin-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/261092>