

Rabbit anti-Human, Pig MAPKAPK3 Polyclonal Antibody | anti-MAPKAPK3 antibody

Anti-MAPKAPK3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA261098	Rabbit	Human, Pig	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

MAPKAPK3; Polyclonal Antibody; Anti-MAPKAPK3 Antibody; MAP kinase-activated protein kinase 3; MAPK-activated protein kinase 3; MAPKAP kinase 3; MAPKAP-K3; MAPKAPK-3; MK-3; Chromosome 3p kinase; 3pK; anti-MAPKAPK3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Pig
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of MAPKAPK3 protein.
Purity/Purification	The antibody was purified by affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	382
Applicable Applications for anti-MAPKAPK3 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human MAPKAPK3. The exact sequence is proprietary.
Preparation and Storage	Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of MAPKAPK3 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



WB (Western Blot)

Western blot analysis of MAPKAPK3 expression in Hela (A), Jurkat (B) whole cell lysates.



IHC (Immunohistochemistry)

Immunohistochemical analysis of MAPKAPK3 staining in human heart formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

ADDITIONAL INFORMATION

Related Product Information for anti-MAPKAPK3 antibody	Rabbit polyclonal antibody to MAPKAPK3
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	74762148
NCBI GeneID	7867
UniProt Accession #	Q16644; B5BU67
NCBI Official Full Name	MAP kinase-activated protein kinase 3
NCBI Official Synonym Full Names	mitogen-activated protein kinase-activated protein kinase 3
NCBI Official Symbol	MAPKAPK3
NCBI Official Synonym Symbols	3PK; MK-3; MAPKAP3; MAPKAP-K3; MAPKAPK-3
NCBI Protein Information	MAP kinase-activated protein kinase 3
UniProt Protein Name	MAP kinase-activated protein kinase 3
UniProt Gene Name	MAPKAPK3
UniProt Synonym Gene Names	MAPK-activated protein kinase 3; MAPKAP kinase 3; MAPKAP-K3; MAPKAPK-3; MK-3; 3pK
UniProt Entry Name	MAPK3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mapkapk3-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/261098>