

Rabbit Alpha-1A Adrenergic Receptor Polyclonal Antibody | anti-ADRA1A antibody

Anti-Alpha-1A Adrenergic Receptor Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA261115	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Alpha-1A Adrenergic Receptor; Polyclonal Antibody; Anti-Alpha-1A Adrenergic Receptor Antibody; ADRA1C; Alpha-1A adrenergic receptor; Alpha-1A adrenoreceptor; Alpha-1A adrenoceptor; Alpha-1C adrenergic receptor; Alpha-adrenergic receptor 1c; anti-ADRA1A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Alpha-1A Adrenergic Receptor protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	466
Applicable Applications for anti-ADRA1A antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Alpha-1A Adrenergic Receptor. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of Alpha-1A Adrenergic Receptor staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Alpha-1A Adrenergic Receptor staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Alpha-1A Adrenergic Receptor expression in LOVO (A), Jurkat (B), NIH3T3 (C), PC12 (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-ADRA1A antibody	Rabbit polyclonal antibody to Alpha-1A Adrenergic Receptor
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	1168246
NCBI GeneID	148
UniProt Accession #	P35348; O60451; Q13675; Q13729; A8K0I3; B0ZBD1; B0ZBD2; B0ZBD4; B0ZBD5; B0ZBD6; B0ZBD8; B0ZBD9
Molecular Weight	40,638 Da
NCBI Official Full Name	Alpha-1A adrenergic receptor
NCBI Official Synonym Full Names	adrenoceptor alpha 1A
NCBI Official Symbol	ADRA1A
NCBI Official Synonym Symbols	ADRA1C; ADRA1L1; ALPHA1AAR
NCBI Protein Information	alpha-1A adrenergic receptor
UniProt Protein Name	Alpha-1A adrenergic receptor
UniProt Gene Name	ADRA1A
UniProt Synonym Gene Names	ADRA1C; Alpha-1A adrenoceptor
UniProt Entry Name	ADA1A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/alpha-1a-adrenergic-receptor-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/261115>