

Rabbit STMN1 Polyclonal Antibody | anti-STMN1 antibody

Anti-STMN1 (pS16) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA261187	Rabbit	Human, Mouse, Rat, Bovine, Chicken, Monkey, Pig	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

STMN1; Polyclonal Antibody; Anti-STMN1 (pS16) Phospho Antibody; C1orf215; LAP18; OP18; Stathmin; Leukemia-associated phosphoprotein p18; Metablastin; Oncoprotein 18; Op18; Phosphoprotein p19; pp19; Proslin; Protein Pr22; pp17; anti-STMN1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Chicken, Monkey, Pig
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of STMN1 (pS16) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	149
Applicable Applications for anti-STMN1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human STMN1 (pS16). The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of STMN1 (pS16) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of STMN1 (pS16) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of STMN1 (pS16) expression in HEK293T (A), HeLa (B), Jurkat (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-STMN1 antibody

Rabbit polyclonal antibody to STMN1 (pS16)

NCBI AND UNIPROT INFORMATION

NCBI GI #	134973
NCBI GeneID	3925
UniProt Accession #	P16949; A2A2D1; B2R4E7; B7Z8N4; D3DPJ5
Molecular Weight	19,824 Da
NCBI Official Full Name	Stathmin
NCBI Official Synonym Full Names	stathmin 1
NCBI Official Symbol	STMN1
NCBI Official Synonym Symbols	Lag; SMN; OP18; PP17; PP19; PR22; LAP18; C1orf215
NCBI Protein Information	stathmin
UniProt Protein Name	Stathmin
UniProt Gene Name	STMN1
UniProt Synonym Gene Names	C1orf215; LAP18; OP18; Op18; pp19
UniProt Entry Name	STMN1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/stmn1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/261187>