

Rabbit Beta-actin Polyclonal Antibody | anti-ACTB antibody

Anti-Beta-actin Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA261215	Rabbit	Human, Mouse, Rat, Rabbit, Chicken, Monkey, Sheep	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Beta-actin; Polyclonal Antibody; Anti-Beta-actin Antibody; Actin cytoplasmic 1; anti-ACTB antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Rabbit, Chicken, Monkey, Sheep
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Beta-actin protein.
Purity/Purification	The antibody was purified by affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	375
Applicable Applications for anti-ACTB antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein corresponding to human Beta-actin.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Beta-actin staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Beta-actin staining in human ovary formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Beta-actin expression in HepG2 (A), 293T (B), mouse kidney (C), rat liver (D), rabbit testis (E), sheep lung (F) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-ACTB antibody	Rabbit polyclonal to Beta-actin
--	---------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	46397333
NCBI GeneID	60
UniProt Accession #	P60709; P02570; P70514; P99021; Q11211; Q64316; Q75MN2; Q96B34; Q96HG5
Molecular Weight	41,737 Da
NCBI Official Full Name	Actin, cytoplasmic 1
NCBI Official Synonym Full Names	actin, beta
NCBI Official Symbol	ACTB
NCBI Official Synonym Symbols	BRWS1; PS1TP5BP1
NCBI Protein Information	actin, cytoplasmic 1
UniProt Protein Name	Actin, cytoplasmic 1
UniProt Gene Name	ACTB
UniProt Entry Name	ACTB_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/beta-actin-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/261215>