

Rabbit GLUR1 Polyclonal Antibody | anti-GRIA1 antibody

Anti-GLUR1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA261406	Rabbit	Human, Mouse, Rat, Monkey	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

GLUR1; Polyclonal Antibody; Anti-GLUR1 Antibody; GLUH1; Glutamate receptor 1; GluR-1; AMPA-selective glutamate receptor 1; GluR-A; GluR-K1; Glutamate receptor ionotropic; AMPA 1; GluA1; anti-GRIA1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of GLUR1 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	906
Applicable Applications for anti-GRIA1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human GLUR1. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of GLUR1 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of GLUR1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of GLUR1 expression in rat kidney (A) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-GRIA1 antibody	Rabbit polyclonal antibody to GLUR1
---	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	116242505
NCBI GeneID	2890
UniProt Accession #	P42261; Q2NKM6; B7Z2S0; B7Z2W8; B7Z3F6; B7Z9G9; D3DQI4; E7ESV8
Molecular Weight	102,864 Da
NCBI Official Full Name	Glutamate receptor 1
NCBI Official Synonym Full Names	glutamate ionotropic receptor AMPA type subunit 1
NCBI Official Symbol	GRIA1
NCBI Official Synonym Symbols	GLUH1; GLUR1; GLURA; GluA1; HBGR1
NCBI Protein Information	glutamate receptor 1
UniProt Protein Name	Glutamate receptor 1
UniProt Gene Name	GRIA1
UniProt Synonym Gene Names	GLUH1; GLUR1; GluR-1; GluA1
UniProt Entry Name	GRIA1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.