

Rabbit SGK1 Polyclonal Antibody | anti-SGK1 antibody

Anti-SGK1 (pS422) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA261439	Rabbit	Human, Monkey	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

SGK1; Polyclonal Antibody; Anti-SGK1 (pS422) Phospho Antibody; SGK; Serine/threonine-protein kinase Sgk1; Serum/glucocorticoid-regulated kinase 1; anti-SGK1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Monkey
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of SGK1 (pS422) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	431
Applicable Applications for anti-SGK1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human SGK1. The exact sequence is proprietary.
Preparation and Storage	Shipped and store at 4 degree C for one year. Do not freeze.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of SGK1 (pS422) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of SGK1 (pS422) staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of SGK1 (pS422) expression in A549 Insulin-treated (A), K562 Insulin-treated (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-SGK1 antibody	Rabbit polyclonal antibody to SGK1 (pS422)
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	90185131
NCBI GeneID	6446
UniProt Accession #	O00141; Q5TCN2; Q5TCN3; Q5TCN4; Q5VY65; Q9UN56; B7UUP7; B7UUP8; B7UUP9; B7Z5B2; E1P583
Molecular Weight	52,119 Da
NCBI Official Full Name	Serine/threonine-protein kinase Sgk1
NCBI Official Synonym Full Names	serum/glucocorticoid regulated kinase 1
NCBI Official Symbol	SGK1
NCBI Official Synonym Symbols	SGK
NCBI Protein Information	serine/threonine-protein kinase Sgk1
UniProt Protein Name	Serine/threonine-protein kinase Sgk1
UniProt Gene Name	SGK1
UniProt Synonym Gene Names	SGK
UniProt Entry Name	SGK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/sgk1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/261439>