

Rabbit MKK3 Polyclonal Antibody | anti-MAP2K3 antibody

Anti-MKK3 (pT222) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA261447	Rabbit	Human, Mouse, Zebrafish	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MKK3; Polyclonal Antibody; Anti-MKK3 (pT222) Phospho Antibody; MEK3; PRKMK3; SKK2; Dual specificity mitogen-activated protein kinase kinase 3; MAP kinase kinase 3; MAPKK 3; MAPK/ERK kinase 3; MEK 3; Stress-activated protein kinase kinase 2; SAPK kinase 2; SAPKK-2; SAPKK2; anti-MAP2K3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Zebrafish
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of MKK3 (pT222) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Sequence Length	347
Applicable Applications for anti-MAP2K3 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human MKK3. The exact sequence is proprietary.
Preparation and Storage	Shipped and store at 4 degree C for one year. Do not freeze.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of MKK3 (pT222) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of MKK3 (pT222) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MKK3 (pT222) expression in HeLa (A), Jurkat (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-MAP2K3 antibody

Rabbit polyclonal antibody to MKK3 (pT222)

NCBI AND UNIPROT INFORMATION

NCBI GI #	24638466
NCBI GeneID	5606
UniProt Accession #	P46734; Q99441; Q9UE71; Q9UE72; B3KSK7
Molecular Weight	39,940 Da
NCBI Official Full Name	Dual specificity mitogen-activated protein kinase kinase 3
NCBI Official Synonym Full Names	mitogen-activated protein kinase kinase 3
NCBI Official Symbol	MAP2K3
NCBI Official Synonym Symbols	MEK3; MKK3; MAPKK3; PRKMK3; SAPKK2; SAPKK-2
NCBI Protein Information	dual specificity mitogen-activated protein kinase kinase 3
UniProt Protein Name	Dual specificity mitogen-activated protein kinase kinase 3
UniProt Gene Name	MAP2K3
UniProt Synonym Gene Names	MEK3; MKK3; PRKMK3; SKK2; MAP kinase kinase 3; MAPKK 3; MEK 3; SAPK kinase 2; SAPKK-2; SAPKK2
UniProt Entry Name	MP2K3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mkk3-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/261447>