

Rabbit RRM2 Polyclonal Antibody | anti-RRM2 antibody

Anti-RRM2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA261821	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

RRM2; Polyclonal Antibody; Anti-RRM2 Antibody; RR2; Ribonucleoside-diphosphate reductase subunit M2; Ribonucleotide reductase small chain; Ribonucleotide reductase small subunit; anti-RRM2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of RRM2 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	389
Applicable Applications for anti-RRM2 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human RRM2
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of RRM2 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of RRM2 staining in rat heart formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of RRM2 expression in MCF7 (A), HeLa (B), Jurakt (C), mouse thymus (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-RRM2 antibody

Rabbit polyclonal antibody to RRM2

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557845
NCBI GeneID	6241
NCBI Accession #	NP_001025.1
NCBI GenBank Nucleotide #	NM_001034.3
UniProt Accession #	P31350; Q5WRU7; B2R9B5; J3KP43
Molecular Weight	51,093 Da
NCBI Official Full Name	ribonucleoside-diphosphate reductase subunit M2 isoform 2
NCBI Official Synonym Full Names	ribonucleotide reductase regulatory subunit M2
NCBI Official Symbol	RRM2
NCBI Official Synonym Symbols	R2; RR2; RR2M
NCBI Protein Information	ribonucleoside-diphosphate reductase subunit M2
UniProt Protein Name	Ribonucleoside-diphosphate reductase subunit M2
UniProt Gene Name	RRM2
UniProt Synonym Gene Names	RR2
UniProt Entry Name	RIR2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/rrm2-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/261821>