

Rabbit DLD Polyclonal Antibody | anti-DLD antibody

Anti-DLD Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA261829	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

DLD; Polyclonal Antibody; Anti-DLD Antibody; GCSL; LAD; PHE3; Dihydrolipoyl dehydrogenase; mitochondrial; Dihydrolipoamide dehydrogenase; Glycine cleavage system L protein; anti-DLD antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of DLD protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	509
Applicable Applications for anti-DLD antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human DLD
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of DLD staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of DLD staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of DLD expression in MCF7 (A), K562 (B), mouse kidney (C), mouse liver (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-DLD antibody	Rabbit polyclonal antibody to DLD
---	-----------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	91199540
NCBI GeneID	1738
NCBI Accession #	NP_000099.2
NCBI GenBank Nucleotide #	NM_000108.4
UniProt Accession #	P09622; Q14131; Q14167; Q59EV8; Q8WTS4; B2R5X0; B4DHG0; B4DT69
Molecular Weight	49,283 Da
NCBI Official Full Name	dihydrolipoyl dehydrogenase, mitochondrial isoform 1
NCBI Official Synonym Full Names	dihydrolipoamide dehydrogenase
NCBI Official Symbol	DLD
NCBI Official Synonym Symbols	E3; LAD; DLDD; DLDH; GCSL; PHE3
NCBI Protein Information	dihydrolipoyl dehydrogenase, mitochondrial
UniProt Protein Name	Dihydrolipoyl dehydrogenase, mitochondrial
UniProt Gene Name	DLD
UniProt Synonym Gene Names	GCSL; LAD; PHE3
UniProt Entry Name	DLDH_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/dld-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/261829>