

Rabbit SerRS Polyclonal Antibody | anti-SARS antibody

Anti-SerRS Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA261852	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

SerRS; Polyclonal Antibody; Anti-SerRS Antibody; SERS; Serine--tRNA ligase; cytoplasmic; Seryl-tRNA synthetase; Seryl-tRNA(Ser/Sec) synthetase; anti-SARS antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of SerRS protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	514
Applicable Applications for anti-SARS antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human SerRS
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of SerRS staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of SerRS staining in human gastric cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of SerRS expression in HepG2 (A), HeLa (B), mouse kidney (C), rat brain (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-SARS antibody

Rabbit polyclonal antibody to SerRS

NCBI AND UNIPROT INFORMATION

NCBI GI #	16306548
NCBI GeneID	6301
NCBI Accession #	NP_006504.2
NCBI GenBank Nucleotide #	NM_006513.3
UniProt Accession #	P49591; Q5T5C8; Q9NSE3; B2R6Y9
Molecular Weight	58,777 Da
NCBI Official Full Name	serine--tRNA ligase, cytoplasmic
NCBI Official Synonym Full Names	seryl-tRNA synthetase
NCBI Official Symbol	SARS
NCBI Official Synonym Symbols	SERS; SERRS
NCBI Protein Information	serine--tRNA ligase, cytoplasmic
UniProt Protein Name	Serine--tRNA ligase, cytoplasmic
UniProt Gene Name	SARS
UniProt Synonym Gene Names	SERS; SerRS
UniProt Entry Name	SYSC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/serrs-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/261852>