

Rabbit PSMB9 Polyclonal Antibody | anti-PSMB9 antibody

Anti-PSMB9 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA262235	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PSMB9; Polyclonal Antibody; Anti-PSMB9 Antibody; LMP2; PSMB6i; RING12; Proteasome subunit beta type-9; Low molecular mass protein 2; Macropain chain 7; Multicatalytic endopeptidase complex chain 7; Proteasome chain 7; Proteasome subunit beta-1i; Really interesting new gene 12 protein; anti-PSMB9 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of PSMB9 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	209
Applicable Applications for anti-PSMB9 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human PSMB9
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of PSMB9 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of PSMB9 staining in mouse kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of PSMB9 expression in HepG2 (A), THP1 (B), mouse liver (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-PSMB9 antibody	Rabbit polyclonal antibody to PSMB9
---	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506205
NCBI GeneID	5698
NCBI Accession #	NP_002791.1
NCBI GenBank Nucleotide #	NM_002800.4
UniProt Accession #	P28065; Q16523; Q5JNW4; B0V0T1
Molecular Weight	22,328 Da
NCBI Official Full Name	proteasome subunit beta type-9
NCBI Official Synonym Full Names	proteasome subunit beta 9
NCBI Official Symbol	PSMB9
NCBI Official Synonym Symbols	LMP2; PSMB6i; RING12; beta1i
NCBI Protein Information	proteasome subunit beta type-9
UniProt Protein Name	Proteasome subunit beta type-9
UniProt Gene Name	PSMB9
UniProt Synonym Gene Names	LMP2; PSMB6i; RING12
UniProt Entry Name	PSB9_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/psmb9-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/262235>