

Rabbit Rac 1 Polyclonal Antibody | anti-RAC1 antibody

Anti-Rac 1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA262312	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Rac 1; Polyclonal Antibody; Anti-Rac 1 Antibody; TC25; Ras-related C3 botulinum toxin substrate 1; Cell migration-inducing gene 5 protein; Ras-like protein TC25; p21-Rac1; anti-RAC1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Rac 1 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	211
Applicable Applications for anti-RAC1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human Rac 1
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Rac 1 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Rac 1 staining in human gastric cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Rac 1 expression in BT474 (A), mouse liver (B), rat brain (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-RAC1 antibody	Rabbit polyclonal antibody to Rac 1
--	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	9845511
NCBI GeneID	5879
NCBI Accession #	NP_008839.2
NCBI GenBank Nucleotide #	NM_006908.4
UniProt Accession #	P63000; O95501; P15154; Q3Y4D3; Q5JAA8; Q9BTB4; AAB22206
Molecular Weight	23,467 Da
NCBI Official Full Name	ras-related C3 botulinum toxin substrate 1 isoform Rac1
NCBI Official Synonym Full Names	ras-related C3 botulinum toxin substrate 1 (rho family, small GTP binding protein Rac1)
NCBI Official Symbol	RAC1
NCBI Official Synonym Symbols	MIG5; Rac-1; TC-25; p21-Rac1
NCBI Protein Information	ras-related C3 botulinum toxin substrate 1
UniProt Protein Name	Ras-related C3 botulinum toxin substrate 1
UniProt Gene Name	RAC1
UniProt Synonym Gene Names	TC25
UniProt Entry Name	RAC1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/rac-1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/262312>