

Rabbit anti-Human HAGH Polyclonal Antibody | anti-HAGH antibody

Anti-HAGH Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA262347	Rabbit	Human	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

HAGH; Polyclonal Antibody; Anti-HAGH Antibody; GLO2; HAGH1; Hydroxyacylglutathione hydrolase; mitochondrial; Glyoxalase II; Glx II; anti-HAGH antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of HAGH protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	236
Applicable Applications for anti-HAGH antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human HAGH
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of HAGH staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of HAGH staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of HAGH expression in HepG2 (A), mouse liver (B), mouse heart (C), rat kidney (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-HAGH antibody	Rabbit polyclonal antibody to HAGH
--	------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	94538320
NCBI GeneID	3029
NCBI Accession #	NP_001035517.1
NCBI GenBank Nucleotide #	NM_001040427.1
UniProt Accession #	Q16775; A8K290; B4DP33; B4DRA7; E7EN93
Molecular Weight	25,455 Da
NCBI Official Full Name	hydroxyacylglutathione hydrolase, mitochondrial isoform 2
NCBI Official Synonym Full Names	hydroxyacylglutathione hydrolase
NCBI Official Symbol	HAGH
NCBI Official Synonym Symbols	GLO2; GLX2; GLXII; HAGH1
NCBI Protein Information	hydroxyacylglutathione hydrolase, mitochondrial
UniProt Protein Name	Hydroxyacylglutathione hydrolase, mitochondrial
UniProt Gene Name	HAGH
UniProt Synonym Gene Names	GLO2; HAGH1; Glx II
UniProt Entry Name	GLO2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/hagh-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/262347>