

Rabbit Ferritin L Polyclonal Antibody | anti-FTL antibody

Anti-Ferritin L Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA262596	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Ferritin L; Polyclonal Antibody; Anti-Ferritin L Antibody; Ferritin light chain; Ferritin L subunit; anti-FTL antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Ferritin L protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	175
Applicable Applications for anti-FTL antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human Ferritin L
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Ferritin L staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Ferritin L staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Ferritin L expression in HEK293T (A), MCF7 (B), HepG2 (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-FTL antibody	Rabbit polyclonal antibody to Ferritin L
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	20149498
NCBI GeneID	2512
NCBI Accession #	NP_000137.2
NCBI GenBank Nucleotide #	NM_000146.3
UniProt Accession #	P02792; Q6IBT7; Q7Z2W1; Q86WI9; Q8WU07; Q96AU9; Q96CU0; Q9BTZ8; B2R4B9
Molecular Weight	20,020 Da
NCBI Official Full Name	ferritin light chain
NCBI Official Synonym Full Names	ferritin light chain
NCBI Official Symbol	FTL
NCBI Official Synonym Symbols	LFTD; NBIA3
NCBI Protein Information	ferritin light chain
UniProt Protein Name	Ferritin light chain
UniProt Gene Name	FTL
UniProt Synonym Gene Names	Ferritin L subunit
UniProt Entry Name	FRIL_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ferritin-l-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/262596>