

Rabbit ERK5 Polyclonal Antibody | anti-MAPK7 antibody

Anti-ERK5 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA262667	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

ERK5; Polyclonal Antibody; Anti-ERK5 Antibody; BMK1; PRKM7; Mitogen-activated protein kinase 7; MAP kinase 7; MAPK 7; Big MAP kinase 1; BMK-1; Extracellular signal-regulated kinase 5; ERK-5; anti-MAPK7 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of ERK5 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	816
Applicable Applications for anti-MAPK7 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human ERK5
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of ERK5 staining in K562 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of ERK5 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of ERK5 expression in K562 (A), Ramos (B), mouse brain (C), mouse lung (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-MAPK7 antibody	Rabbit polyclonal antibody to ERK5
---	------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	20986497
NCBI GeneID	5598
NCBI Accession #	NP_002740.2
NCBI GenBank Nucleotide #	NM_002749.3
UniProt Accession #	Q13164; Q16634; Q59F50; Q6QLU7; Q7L4P4; Q969G1; Q96G51
Molecular Weight	50,152 Da
NCBI Official Full Name	mitogen-activated protein kinase 7 isoform 1
NCBI Official Synonym Full Names	mitogen-activated protein kinase 7
NCBI Official Symbol	MAPK7
NCBI Official Synonym Symbols	BMK1; ERK4; ERK5; PRKM7
NCBI Protein Information	mitogen-activated protein kinase 7
UniProt Protein Name	Mitogen-activated protein kinase 7
UniProt Gene Name	MAPK7
UniProt Synonym Gene Names	BMK1; ERK5; PRKM7; MAP kinase 7; MAPK 7; BMK-1; ERK-5
UniProt Entry Name	MK07_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/erk5-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/262667>