

Rabbit NDRG1 Polyclonal Antibody | anti-NDRG1 antibody

Anti-NDRG1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA262689	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

NDRG1; Polyclonal Antibody; Anti-NDRG1 Antibody; CAP43; DRG1; RTP; Protein NDRG1; Differentiation-related gene 1 protein; DRG-1; N-myc downstream-regulated gene 1 protein; Nickel-specific induction protein Cap43; Reducing agents and tunicamycin-responsive protein; Rit42; anti-NDRG1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of NDRG1 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	394
Applicable Applications for anti-NDRG1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human NDRG1
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of NDRG1 staining in SW480 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of NDRG1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of NDRG1 expression in Hela (A), SW480 (B), mouse kidney (C), mouse brain (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-NDRG1 antibody	Rabbit polyclonal antibody to NDRG1
---	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	207028748
NCBI GeneID	10397
NCBI Accession #	NP_001128714.1
NCBI GenBank Nucleotide #	NM_001135242.1
UniProt Accession #	Q92597; O15207; Q6IBG2; Q9NYR6; Q9UK29; B3KR80; B7Z446
Molecular Weight	33,650 Da
NCBI Official Full Name	protein NDRG1 isoform 1
NCBI Official Synonym Full Names	N-myc downstream regulated 1
NCBI Official Symbol	NDRG1
NCBI Official Synonym Symbols	GC4; RTP; DRG1; NDR1; NMSL; TDD5; CAP43; CMT4D; DRG-1; HMSNL; RIT42; TARG1; PROXY1
NCBI Protein Information	protein NDRG1
UniProt Protein Name	Protein NDRG1
UniProt Gene Name	NDRG1
UniProt Synonym Gene Names	CAP43; DRG1; RTP; DRG-1; RTP
UniProt Entry Name	NDRG1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ndrg1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/262689>