

Rabbit NFAT2 Polyclonal Antibody | anti-NFATC1 antibody

Anti-NFAT2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA262694	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

NFAT2; Polyclonal Antibody; Anti-NFAT2 Antibody; NFATC; Nuclear factor of activated T-cells; cytoplasmic 1; NF-ATc1; NFATc1; NFAT transcription complex cytosolic component; NF-ATc; NFATc; anti-NFATC1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of NFAT2 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	943
Applicable Applications for anti-NFATC1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human NFAT2
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of NFAT2 staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of NFAT2 staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of NFAT2 expression in Jurkat (A), HL60 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-NFATC1 antibody	Rabbit polyclonal antibody to NFAT2
--	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	522021835
NCBI GeneID	4772
NCBI Accession #	NP_001265598.1
NCBI GenBank Nucleotide #	NM_001278669.1
UniProt Accession #	O95644; Q12865; Q15793; Q2M1S3; B5B2M4; B5B2M5; B5B2M6; B5B2M7; B5B2M8; B5B2M9; B5B2N1
Molecular Weight	38,626 Da
NCBI Official Full Name	nuclear factor of activated T-cells, cytoplasmic 1 isoform F
NCBI Official Synonym Full Names	nuclear factor of activated T-cells 1
NCBI Official Symbol	NFATC1
NCBI Official Synonym Symbols	NFAT2; NFATc; NF-ATC; NF-ATc1.2
NCBI Protein Information	nuclear factor of activated T-cells, cytoplasmic 1
UniProt Protein Name	Nuclear factor of activated T-cells, cytoplasmic 1
UniProt Gene Name	NFATC1
UniProt Synonym Gene Names	NFAT2; NFATC; NF-ATc1; NFATc1; NF-ATc; NFATc
UniProt Entry Name	NFAC1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/nfat2-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/262694>