

Rabbit LRH-1 Polyclonal Antibody | anti-NR5A2 antibody

Anti-LRH-1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA262700	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

LRH-1; Polyclonal Antibody; Anti-LRH-1 Antibody; B1F; CPF; FTF; Nuclear receptor subfamily 5 group A member 2; Alpha-1-fetoprotein transcription factor; B1-binding factor; hB1F; CYP7A promoter-binding factor; Hepatocytic transcription factor; Liver receptor homolog 1; anti-NR5A2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of LRH-1 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	469
Applicable Applications for anti-NR5A2 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human LRH-1
Preparation and Storage	Upon receipt, aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of LRH-1 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of LRH-1 staining in mouse lung formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of LRH-1 expression in MCF7 (A), Hela (B), mouse liver (C), rat liver (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-NR5A2 antibody	Rabbit polyclonal antibody to LRH-1
---	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	451172094
NCBI GeneID	2494
NCBI Accession #	NP_001263393.1
NCBI GenBank Nucleotide #	NM_001276464.1
UniProt Accession #	O00482; O95642; Q147U3; B4E2P3
NCBI Official Full Name	nuclear receptor subfamily 5 group A member 2 isoform 3
NCBI Official Synonym Full Names	nuclear receptor subfamily 5 group A member 2
NCBI Official Symbol	NR5A2
NCBI Official Synonym Symbols	B1F; CPF; FTF; B1F2; LRH1; LRH-1; FTZ-F1; hB1F-2; FTZ-F1beta
NCBI Protein Information	nuclear receptor subfamily 5 group A member 2
UniProt Protein Name	Nuclear receptor subfamily 5 group A member 2
UniProt Gene Name	NR5A2
UniProt Synonym Gene Names	B1F; CPF; FTF; hB1F; LRH-1
UniProt Entry Name	NR5A2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/lrh-1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/262700>