

Rabbit SMAD2 Polyclonal Antibody | anti-SMAD2 antibody

Anti-SMAD2 (pT220) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA263987	Rabbit	Human, Mouse, Rat, Bovine, Chicken, Zebrafish	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

SMAD2; Polyclonal Antibody; Anti-SMAD2 (pT220) Antibody; MADH2; MADR2; Mothers against decapentaplegic homolog 2; MAD homolog 2; Mothers against DPP homolog 2; JV18-1; Mad-related protein 2; hMAD-2; SMAD family member 2; SMAD 2; Smad2; hSMAD2; anti-SMAD2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Chicken, Zebrafish
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of SMAD2 (pT220) protein.
Purity/Purification	Purified by Immunogen Affinity Chromatography
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Applicable Applications for anti-SMAD2 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human SMAD2 (pT220). The exact sequence is proprietary.
Preparation and Storage	Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of SMAD2 (pT220) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 degree C in a humidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of SMAD2 (pT220) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of SMAD2 (pT220) expression in rat heart (A) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-SMAD2 antibody	Rabbit polyclonal antibody to SMAD2
---	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	51173730
NCBI GeneID	4087
NCBI Accession #	NP_001003652.1
NCBI GenBank Nucleotide #	NM_001003652.3
UniProt Accession #	Q15796; Q62432; O70436
Molecular Weight	52,306 Da
NCBI Official Full Name	mothers against decapentaplegic homolog 2 isoform 1
NCBI Official Synonym Full Names	SMAD family member 2
NCBI Official Symbol	SMAD2
NCBI Official Synonym Symbols	JV18; MADH2; MADR2; JV18-1; hMAD-2; hSMAD2
NCBI Protein Information	mothers against decapentaplegic homolog 2; MAD homolog 2; mother against DPP homolog 2; Sma- and Mad-related protein 2; SMAD, mothers against DPP homolog 2
UniProt Protein Name	Mothers against decapentaplegic homolog 2
UniProt Gene Name	SMAD2
UniProt Synonym Gene Names	MADH2; MADR2; MAD homolog 2; Mothers against DPP homolog 2; hMAD-2; SMAD 2; Smad2; hSMAD2
UniProt Entry Name	SMAD2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/smad2-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/263987>