

Rabbit CD124 Polyclonal Antibody | anti-CD124 antibody

Anti-CD124 (pY497) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA263992	Rabbit	Human, Mouse, Rat, Pig	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

CD124; Polyclonal Antibody; Anti-CD124 (pY497) Antibody; IL4RA; Interleukin-4 receptor subunit alpha; IL-4 receptor subunit alpha; IL-4R subunit alpha; IL-4R-alpha; IL-4RA; anti-CD124 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Pig
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of CD124 (pY497) protein.
Purity/Purification	Purified by Immunogen Affinity Chromatography
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Applicable Applications for anti-CD124 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CD124 (pY497). The exact sequence is proprietary.
Preparation and Storage	Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of CD124 (pY497) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 degree C in a humidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of CD124 (pY497) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of CD124 (pY497) expression in Myla2059 (A), SGC7901 (B), mouse testis (C), rat testis (D) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-CD124 antibody	Rabbit polyclonal antibody to CD124
---	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557669
NCBI GeneID	3566
NCBI Accession #	NP_000409.1
NCBI GenBank Nucleotide #	NM_000418.3
UniProt Accession #	P24394; P16382; Q63257
Molecular Weight	89,658 Da
NCBI Official Full Name	interleukin-4 receptor subunit alpha isoform a
NCBI Official Synonym Full Names	interleukin 4 receptor
NCBI Official Symbol	IL4R
NCBI Official Synonym Symbols	CD124; IL4RA; IL-4RA
NCBI Protein Information	interleukin-4 receptor subunit alpha; IL4R nirs variant 1; IL-4 receptor subunit alpha; interleukin-4 receptor alpha chain
UniProt Protein Name	Interleukin-4 receptor subunit alpha
UniProt Gene Name	IL4R
UniProt Synonym Gene Names	IL4RA; IL-4 receptor subunit alpha; IL-4R subunit alpha; IL-4R-alpha; IL-4RA; Soluble IL-4 receptor subunit alpha; Soluble IL-4R-alpha; sIL4Ralpha/prot; IL4-BP
UniProt Entry Name	IL4RA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd124-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/263992>