

Rabbit Aconitase 1 Polyclonal Antibody | anti-ACO1 antibody

Anti-Aconitase 1 (pS138) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA263994	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Aconitase 1; Polyclonal Antibody; Anti-Aconitase 1 (pS138) Antibody; IREB1; Cytoplasmic aconitate hydratase; Aconitase; Citrate hydro-lyase; Ferritin repressor protein; Iron regulatory protein 1; IRP1; Iron-responsive element-binding protein 1; IRE-BP 1; anti-ACO1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Aconitase 1 (pS138) protein.
Purity/Purification	Purified by Immunogen Affinity Chromatography
Form/Format	Liquid in 0.42% Potassium Phosphate, 0.87% Sodium Chloride, pH 7.3, 30% Glycerol, and 0.01% Sodium Azide.
Applicable Applications for anti-ACO1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Aconitase 1 (pS138). The exact sequence is proprietary.
Preparation and Storage	Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Aconitase 1 (pS138) staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 degree C in a humidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Aconitase 1 (pS138) staining in human ovary formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Aconitase 1 (pS138) expression in HeLa (A) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-ACO1 antibody

Rabbit polyclonal antibody to Aconitase 1

NCBI AND UNIPROT INFORMATION

NCBI GI #	8659555
NCBI GeneID	48
NCBI Accession #	NP_002188.1
NCBI GenBank Nucleotide #	NM_002197.2
UniProt Accession #	P21399; P28271; Q63270
Molecular Weight	98,399 Da
NCBI Official Full Name	cytoplasmic aconitate hydratase
NCBI Official Synonym Full Names	aconitase 1, soluble
NCBI Official Symbol	ACO1
NCBI Official Synonym Symbols	IRP1; ACONS; IREB1; IREBP; IREBP1
NCBI Protein Information	cytoplasmic aconitate hydratase; IRE-BP 1; citrate hydro-lyase; iron regulatory protein 1; ferritin repressor protein; aconitate hydratase, cytoplasmic; iron-responsive element binding protein 1; iron-responsive element-binding protein 1
UniProt Protein Name	Cytoplasmic aconitate hydratase
UniProt Gene Name	ACO1
UniProt Synonym Gene Names	IREB1; Aconitase; IRP1; IRE-BP 1
UniProt Entry Name	ACOC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/aconitase-1-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/263994>