Mouse FGR Monoclonal Antibody | anti-FGR antibody

FGR (Gardner-Rasheed feline Sarcoma Viral (v-fgr) Oncogene homolog, FLJ43153, MGC75096, SRC2, c-fgr, c-src2, p55c-fgr, p58c-fgr) (HRP)

CATALOG #	HOST	APPLICATIONS
AAA26424	**Mouse**	**Immunohistochemistry, Western Blot**

PRODUCT SYNONYMS

FGR; Monoclonal Antibody; FGR (Gardner-Rasheed feline Sarcoma Viral (v-fgr) Oncogene homolog; FLJ43153; MGC75096; SRC2; c-fgr; c-src2; p55c-fgr; p58c-fgr) (HRP); Gardner-Rasheed feline Sarcoma Viral (v-fgr) Oncogene homolog; p58c-fgr; anti-FGR antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a,k
Clone Number	3G10
Specificity	Recognizes FGR.
Purity/Purification	Purified
Form/Format	Supplied as a liquid in PBS, pH 7.2. No preservative added. Labeled with horseradish peroxidase (HRP).
Sequence Length	529
Applicable Applications for anti-FGR antibody	IHC (Immunohistochemistry), WB (Western Blot)
Application Notes	Applications are based on unconjugated antibody.
Immunogen	FGR (AAH64382, 1aa-90aa) partial recombinant protein with GST tag. MW of the GST tag alone is 26kD.
Immunogen Sequence	MGCVFCKKLEPVATAKEDAGLEGDFRSYGAADHYGPDPTKARPASSFAHIPNYSNFSSQAINPGFLDSGTIRGVSGIGVTLFIALYDYEA
Conjugate	HRP
Preparation and Storage	Store product at 4 degree C if to be used immediately within two weeks. For long-term storage, aliquot to avoid repeated freezing and thawing and store at -20 degree C. Aliquots are stable at -20 degree C for 12 months after receipt. Dilute required amount only prior to immediate use. Further dilutions can be made in assay buffer. Note: Sodium azide is a potent inhibitor of peroxidase and should not be added to HRP conjugates. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap.

IMAGES

WB (Western Blot)
FGR monoclonal antibody (M01), clone 3G10. Western Blot analysis of FGR expression in Raw 264.7 (Cat # L024V1).

WB (Western Blot)
FGR monoclonal antibody (M01), clone 3G10. Western Blot analysis of FGR expression in PC-12 (Cat # L012V1).

Application Data
Detection limit for recombinant GST tagged FGR is approximately 0.3ng/ml as a capture antibody.

WB (Western Blot)
FGR monoclonal antibody (M01), clone 3G10 Western Blot analysis of FGR expression in HeLa (Cat # L013V1).

IHC (Immunohistochemistry)
Immunoperoxidase of monoclonal antibody to FGR on formalin-fixed paraffin-embedded human lymph node. [antibody concentration 3 ug/ml]

Application Data
Immunoperoxidase of monoclonal antibody to FGR on formalin-fixed paraffin-embedded human lymph node. [antibody concentration 3 ug/ml]

ADDITIONAL INFORMATION

Product Categories/Family for anti-FGR antibody	Antibodies; Protein Kinases

NCBI AND UNIPROT INFORMATION

NCBI GI #	39963676
NCBI GeneID	2268
NCBI Official Full Name	Gardner-Rasheed feline sarcoma viral (v-fgr) oncogene homolog
NCBI Official Synonym Full Names	FGR proto-oncogene, Src family tyrosine kinase
NCBI Official Symbol	FGR
NCBI Official Synonym Symbols	SRC2; c-fgr; c-src2; p55-Fgr; p58-Fgr; p55c-fgr; p58c-fgr
NCBI Protein Information	tyrosine-protein kinase Fgr

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/monoclonal/fgr-antibody-clone-3g10-immunohistochemistry-western-blot-elisa/26424