

Mouse anti-Human MUC1 Monoclonal Antibody | anti-MUC1 antibody

Anti-MUC1 Antibody

CATALOG # AAA264829	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

MUC1; Monoclonal Antibody; Anti-MUC1 Antibody; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Cancer antigen 15-3; CA 15-3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; Polymorphic epithelial mucin; PEM; Tumor-associated epithelial membrane antigen; EMA; Tumor-associated mucin; CD227; anti-MUC1 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG2b
Specificity	Recognizes endogenous levels of MUC1 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in PBS containing 50% glycerol, 0.2% BSA and 0.01% sodium azide.
Applicable Applications for anti-MUC1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within human MUC1. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunohistochemical analysis of MUC1 staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



IHC (Immunohistochemistry)

Immunohistochemical analysis of MUC1 staining in human breast carcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

ADDITIONAL INFORMATION

Related Product Information for anti-MUC1 antibody	Mouse monoclonal antibody to MUC1
--	-----------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	67189007
-----------	----------

NCBI GeneID	4582
NCBI Accession #	NP_001018016.1
NCBI GenBank Nucleotide #	NM_001018016.2
UniProt Accession #	P15941 ; P13931 ; P15942 ; P17626 ; Q0VAP6 ; Q14128 ; Q14876 ; Q16437 ; Q16442 ; Q16615 ; B1AVQ8 ; B1AVR0
Molecular Weight	122,102 Da
NCBI Official Full Name	mucin-1 isoform 2
NCBI Official Synonym Full Names	mucin 1, cell surface associated
NCBI Official Symbol	MUC1
NCBI Official Synonym Symbols	EMA; PEM; PUM; KL-6; MAM6; PEMT; CD227; H23AG; MUC-1; CA 15-3; MUC-1/X; MUC1/ZD; MUC-1/SEC
NCBI Protein Information	mucin-1; episialin; DF3 antigen; H23 antigen; cancer antigen 15-3; krebs von den Lungen-6; mucin 1, transmembrane; tumor-associated mucin; carcinoma-associated mucin; polymorphic epithelial mucin; peanut-reactive urinary mucin; tumor associated epithelial
UniProt Protein Name	Mucin-1
UniProt Gene Name	MUC1
UniProt Synonym Gene Names	PUM; MUC-1; CA 15-3; KL-6; PUM; PEM; EMA; MUC1-NT; MUC1-alpha; MUC1-beta
UniProt Entry Name	MUC1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/muc1-antibody-immunohistochemistry-western-blot-elisa/264829>