

Mouse anti-Human Cytokeratin 13 Monoclonal Antibody | anti-KRT13 antibody

Anti-Cytokeratin 13 Antibody

CATALOG # AAA264861	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

Cytokeratin 13; Monoclonal Antibody; Anti-Cytokeratin 13 Antibody; Keratin type I cytoskeletal 13; Cytokeratin-13; CK-13; Keratin-13; K13; anti-KRT13 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG1
Specificity	Recognizes endogenous levels of Cytokeratin 13 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in PBS containing 50% glycerol, 0.2% BSA and 0.01% sodium azide.
Applicable Applications for anti-KRT13 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within human Cytokeratin 13. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IP (Immunoprecipitation)

Immunohistochemical analysis of Cytokeratin 13 staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Cytokeratin 13 staining in human transitional cell carcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



IF (Immunofluorescence)

Immunohistochemical analysis of Cytokeratin 13 staining in human squamous cell lung carcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

ADDITIONAL INFORMATION

Related Product Information for anti-KRT13 antibody	Mouse monoclonal antibody to Cytokeratin 13
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	30377
NCBI GeneID	3860
UniProt Accession #	P13646; Q53G54; Q6AZK5; Q8N240
Molecular Weight	38,581 Da
NCBI Official Full Name	cytokeratin 13
NCBI Official Synonym Full Names	keratin 13
NCBI Official Symbol	KRT13
NCBI Official Synonym Symbols	K13; CK13; WSN2
NCBI Protein Information	keratin, type I cytoskeletal 13
UniProt Protein Name	Keratin, type I cytoskeletal 13
UniProt Gene Name	KRT13
UniProt Synonym Gene Names	CK-13; K13

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cytokeratin-13-antibody-immunohistochemistry-western-blot-elisa/264861>