

Mouse anti-Human Claudin 7 Monoclonal Antibody | anti-CLDN7 antibody

Anti-Claudin 7 Antibody

CATALOG # AAA264916	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

Claudin 7; Monoclonal Antibody; Anti-Claudin 7 Antibody; CEPTRL2; CPETRL2; Claudin-7; CLDN-7; anti-CLDN7 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG2a
Specificity	Recognizes endogenous levels of Claudin 7 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in PBS containing 50% glycerol, 0.2% BSA and 0.01% sodium azide.
Applicable Applications for anti-CLDN7 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within human Claudin 7. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IP (Immunoprecipitation)

Immunohistochemical analysis of Claudin 7 staining in human kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.




IF (Immunofluorescence)

Immunohistochemical analysis of Claudin 7 staining in human colon carcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Claudin 7 staining in human chromophobe cell renal carcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Claudin 7 expression in LNCaP (A) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-CLDN7 antibody

Mouse monoclonal antibody to Claudin 7

NCBI AND UNIPROT INFORMATION

NCBI GI #	297206811
NCBI GeneID	1366
NCBI Accession #	NP_001171951.1
NCBI GenBank Nucleotide #	NM_001185022.1
UniProt Accession #	O95471; Q6IPN3; Q7Z4Y7; Q9BVN0; B2R9X7; D3DTP0
Molecular Weight	22,418 Da
NCBI Official Full Name	claudin-7 isoform 1
NCBI Official Synonym Full Names	claudin 7
NCBI Official Symbol	CLDN7
NCBI Official Synonym Symbols	CLDN-7; CEPTRL2; CPETRL2; Hs.84359; claudin-1
NCBI Protein Information	claudin-7; clostridium perfringens enterotoxin receptor-like 2
UniProt Protein Name	Claudin-7
UniProt Gene Name	CLDN7
UniProt Synonym Gene Names	CEPTRL2; CPETRL2; CLDN-7
UniProt Entry Name	CLD7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/claudin-7-antibody-immunohistochemistry-western-blot-elisa/264916>