

Mouse anti-Human CD3e Monoclonal Antibody | anti-CD3E antibody

Anti-CD3e Antibody

CATALOG # AAA264945	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

CD3e; Monoclonal Antibody; Anti-CD3e Antibody; T3E; T-cell surface glycoprotein CD3 epsilon chain; T-cell surface antigen T3/Leu-4 epsilon chain; anti-CD3E antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG2a
Specificity	Recognizes endogenous levels of CD3e protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in PBS containing 50% glycerol, 0.2% BSA and 0.01% sodium azide.
Applicable Applications for anti-CD3E antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within human CD3e. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IP (Immunoprecipitation)

Immunohistochemical analysis of CD3e staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



IF (Immunofluorescence)

Immunohistochemical analysis of CD3e staining in human spleen formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



IHC (Immunohistochemistry)

Immunohistochemical analysis of CD3e staining in human diffuse large B-cell lymphoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

ADDITIONAL INFORMATION

Related Product Information for anti-CD3E antibody	Mouse monoclonal antibody to CD3e
--	-----------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	1345708
NCBI GeneID	916
UniProt Accession #	P07766; A8K997
Molecular Weight	207
NCBI Official Full Name	T-cell surface glycoprotein CD3 epsilon chain
NCBI Official Synonym Full Names	CD3e molecule, epsilon (CD3-TCR complex)
NCBI Official Symbol	CD3E
NCBI Official Synonym Symbols	T3E; TCRE; IMD18
NCBI Protein Information	T-cell surface glycoprotein CD3 epsilon chain; CD3-epsilon; T-cell surface antigen T3/Leu-4 epsilon chain; CD3e antigen, epsilon polypeptide (TiT3 complex); T-cell antigen receptor complex, epsilon subunit of T3
UniProt Protein Name	T-cell surface glycoprotein CD3 epsilon chain
UniProt Gene Name	CD3E
UniProt Synonym Gene Names	T3E
UniProt Entry Name	CD3E_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd3e-antibody-immunohistochemistry-western-blot-elisa/264945>